



Conscious behavior after traumatic brain injury: Anatomo-functional support and therapeutic prospects

Jean-Jacques Lemaire, Bénédicte Pontier, Jerome Coste, Thierry Gillart,
Laurent Sakka, Fabien Feschet, Jean Gabrillargues, Emmanuel Coudeyre,
Jacques Luauté

► To cite this version:

Jean-Jacques Lemaire, Bénédicte Pontier, Jerome Coste, Thierry Gillart, Laurent Sakka, et al.. Conscious behavior after traumatic brain injury: Anatomo-functional support and therapeutic prospects. 26ème Congrès de la SOFMER, Société Française de Médecine Physique et de Réadaptation (SOFMER), Oct 2011, Nantes, France. pp.e213-e214, 10.1016/j.rehab.2011.07.387 . hal-01871747

HAL Id: hal-01871747

<https://uca.hal.science/hal-01871747>

Submitted on 26 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Conscious behavior after traumatic brain injury: Anatomo-functional support and therapeutic prospects

J.-J.Lemaire^{a,*}, B.Pontier^b, J.Coste^a, T.Gillart^c, L.Sakka^a, F.Feschet^d,
J.Gabrillargues^e, E.Coudeyre^f, J.Luauté^g

^a Service de neurochirurgie, IGCNC, université d'Auvergne, CHU de Clermont-Ferrand, 63000 Clermont-Ferrand, France

^b Service de MPR, IGCNC, université d'Auvergne, CHU de Clermont-Ferrand, Clermont-Ferrand, France

^c IGCNC, département d'anesthésie-réanimation, université d'Auvergne, CHU de Clermont-Ferrand, Clermont-Ferrand, France

^d IGCNC, université d'Auvergne, Clermont-Ferrand, France

^e Unité de neuroradiologie, IGCNC, université d'Auvergne, CHU de Clermont-Ferrand, Clermont-Ferrand, France

^f Service de MPR, CHU de Clermont-Ferrand, Clermont-Ferrand, France

^g Service de MPR, hospices civils de Lyon, Lyon, France

^{*} speaker

Keywords: Coma, Consciousness, Tegmentum, Thalamus, Basal ganglia, Default-mode network, Precuneus

Objective. Most brain-injured patients with severe and chronic consciousness disorders are in a therapeutic deadlock. This concerns mainly vegetative or neurovegetative patients, and patients in minimally conscious state. Chronic coma is an exceptional condition; certain conditions of akinetic mutism, which are more frequent, can be included in severe and chronic consciousness disorders. The goal is to review the functional connectivity of conscious behaviours and relational arousal, in particular since the introduction of modern clinical imagery.

Description. The connectivity described in this work relies mainly on two magnetic resonance imaging structural studies of the deep brain: a high-resolution atlas (voxel = 250 μ m side; 4.7-Tesla) of an human anatomic piece; an extensive study of deep fascicles (diffusion tensor imaging and tractography; voxel = 1.25 $\times$ 1.25 $\times$ 1.5 mm³; 3-Tesla) on 6 healthy subjects. The results show the support of the functional connectivity of consciousness that involves the mesencephalo-pontine tegmentum, the basal ganglia, the hypothalamus and the thalamus. These deep located regions are connected with the cortex through three main paths: thalamic, ganglionic and rostroventral. The thalamic path rises from the tegmentum, uses the central tegmental tract, and reaches the reticular and dorsomedial thalamus; from the thalamus it spreads to the cortex, the limbic system, the striatum and the pallidum. The ganglionic path uses the lenticular nucleus and projects to the cortex. The rostro-ventral path goes through the ventral tegmental area (below the thalamus) and the posterolateral hypothalamus, and then reaches the frontobasal region; this path uses the basal forebrain bundle.

Prospective. The knowledge of structures controlling conscious behaviours can enable to better understand different types of severe and chronic consciousness disorders. This also could allow proposing adjusted therapeutic options including physical medicine, rehabilitation, pharmacology and neuromodulation.